

## Literaturliste

zum Beitrag „Kieferorthopädie in Deutschland: aggressives Marketing und ineffiziente Behandlung“  
von Dr. Henning Maden, Kieferorthopädie

Forum für Zahnheilkunde 121, Dezember 2014, Seite 6-12

1. Hafez HS, Shaarawy SM, Al-Sakiti AA, Mostafa YA. Dental crowding as a caries risk factor: a systematic review. *Am J Orthod Dentofacial Orthop* 2012; 142(4):443-50.
2. Ainamo J. Relationship between malalignment of the teeth and periodontal disease. *Scand J Dent Res* 1972;80(2):104-10.
3. Buckley LA. The relationships between malocclusion, gingival inflammation, plaque and calculus. *J Periodontol* 1981;52(1):35-40.
4. Bollen AM, Cunha-Cruz J, Bakko DW, Huang GJ, Hujuel PP. The effects of orthodontic therapy on periodontal health: a systematic review of controlled evidence. *J Am Dent Assoc* 2008;139(4):413-22.
5. Huang G. Fourth and forty? *Am J Orthod Dentofacial Orthop* 2008;134(2):174-5.
6. Gesch D, Bernhardt O, Kocher T, et al. Association of malocclusion and functional occlusion with signs of temporomandibular disorders in adults: results of the population-based study of health in Pomerania. *Angle Orthod* 2004;74(4):512-20.
7. Egermark I, Magnusson T, Carlsson GE. A 20-year follow-up of signs and symptoms of temporomandibular disorders and malocclusions in subjects with and without orthodontic treatment in childhood. *Angle Orthod* 2003;73(2):109-15.
8. McNamara JA, Jr., Turp JC. Orthodontic treatment and temporomandibular disorders: is there a relationship? Part 1: Clinical studies. *Fortschr Kieferorthop/J Orofac Orthop* 1997;58(2):74-89.
9. Mohlin B, Axelsson S, Paulin G, et al. TMD in relation to malocclusion and orthodontic treatment. *Angle Orthod* 2007;77(3):542-8.
10. Michelotti A, Iodice G. The role of orthodontics in temporomandibular disorders. *J Oral Rehabil* 2010;37(6):411-29.
11. Shulman JD, Peterson J. The association between incisor trauma and occlusal characteristics in individuals 8-50 years of age. *Dent Traumatol* 2004;20(2):67-74.
12. Koroluk LD, Tulloch JF, Phillips C. Incisor trauma and early treatment for Class II Division 1 malocclusion. *Am J Orthod Dentofacial Orthop* 2003;123(2):117-25; discussion 25-6.
13. English JD, Buschang PH, Throckmorton GS. Does malocclusion affect masticatory performance? *Angle Orthod* 2002;72(1):21-7.
14. Farronato G, Giannini L, Riva R, Galbiati G, Maspero C. Correlations between malocclusions and dyslalias. *Eur J Paediatr Dent* 2012;13(1):13-8.
15. Warren WW, Spalding PM. Dentofacial morphology and oral breathing: a century of controversy. In: Melsen B, editor. *Current controversies in orthodontics*. Chicago: Quintessence Publishing Co; 1991. p. 45-76.
16. Flores-Mir C, Korayem M, Heo G, et al. Craniofacial morphological characteristics in children with obstructive sleep apnea syndrome: a systematic review and meta-analysis. *J Am Dent Assoc* 2013;144(3):269-77.
17. Hanggi MP, Teuscher UM, Roos M, Peltomaki TA. Long-term changes in pharyngeal airway dimensions following activator-headgear and fixed appliance treatment. *Eur J Orthod* 2008;30(6):598-605.
18. Schutz TC, Dominguez GC, Hallinan MP, Cunha TC, Tufik S. Class II correction improves nocturnal breathing in adolescents. *Angle Orthod* 2011;81(2):222-8.
19. Godt A, Koos B, Hagen H, Goz G. Changes in upper airway width associated with Class II treatments (headgear vs activator) and different growth patterns. *Angle Orthod* 2011;81(3):440-6.
20. Altalibi M, Saltaji H, Roduta Roberts M, et al. Developing an index for the orthodontic treatment need in paediatric patients with obstructive sleep apnoea: a protocol for a novel communication tool between physicians and orthodontists. *BMJ Open* 2014;4(9):e005680.
21. Manfredini D, Castroflorio T, Perinetti G, Guarda-Nardini L. Dental occlusion, body posture and temporomandibular disorders: where we are now and where we are heading for. *J Oral Rehabil* 2012;39(6):463-71.
22. Zhou Y, Wang Y, Wang X, Voliere G, Hu R. The impact of orthodontic treatment on the quality of life a systematic review. *BMC Oral Health* 2014;14:66.
23. O'Brien K, Wright J, Conboy F, et al. Effectiveness of early orthodontic treatment with the Twin-block appliance: a multicenter, randomized, controlled trial. Part 2: Psychosocial effects. *Am J Orthod Dentofacial Orthop* 2003;124(5):488-94; discussion 94-5.
24. Feu D, Miguel JA, Celeste RK, Oliveira BH. Effect of orthodontic treatment on oral health-related quality of life. *Angle Orthod* 2013;83(5):892-8.
25. Mohlin B, Kurol J. To what extent do deviations from an ideal occlusion constitute a health risk? *Swed Dent J* 2003;27(1):1-10.
26. Kean MR, Ackerman JL, Vig PS. Malocclusion: beyond the Wendell L.Wylie legacy. *Angle Orthod* 2009;79(1):200-1.
27. Pichelmayr M. Die Rolle des Pädiaters in der Früherkennung von Bissfehlstellungen. *Kieferorthopädische Behandlungen im Überblick*. Pädiatrische Praxis 2005;66:481-96.
28. Ackerman MB. Selling orthodontic need: innocent business decision or guilty pleasure? *Journal of medical ethics* 2010;36(5):275-8.
29. Krey KF, Hirsch C. Frequency of orthodontic treatment in German children and adolescents: influence of age, gender, and socio-economic status.

30. Eur J Orthod 2012;34(2):152-7.
31. Bauer J, Neumann T, Saekel R. Kieferorthopädische Leistungen. In: Zahnmedizinische Versorgung in Deutschland. Bern: Verlag Hans Huber; 2009.
32. Richmond S, Shaw WC, O'Brien KD, et al. The development of the PAR Index (Peer Assessment Rating): reliability and validity. Eur J Orthod 1992;14(2):125-39.
33. Bellot-Arcis C, Montiel-Company JM, Almerich-Silla JM, Paredes-Gallardo V, Gandia-Franco JL. The use of occlusal indices in high-impact literature. Community dental health 2012;29(1):45-8.
34. Aull S. Qualitätssicherung in der kieferorthopädischen Praxis unter Verwendung des PAR Index [Med. Diss.]. Tübingen: Eberhard-Karls-Universität Tübingen; 2001.
35. Fellner U, Schlömer R. der PAR-Index – eine praktikable Methode der Qualitätssicherung. Kieferorthop 1996;10:193-200.
36. Mayes S. Odontometrische Langzeitstudie zur Stabilität kieferorthopädischer Behandlungsergebnisse anhand des PAR-Index [Med. Diss.]. Freiburg: Albert-Ludwigs-Universität Freiburg i. Br.; 1997.
37. Schmidt-Rogge N. Quantitative und qualitative Auswertung kieferorthopädischer Behandlungen [Med. Diss.]. Berlin: Humboldt-Universität zu Berlin; 2001.
38. Hetz T. Vergleichende retrospektive Bewertung des Therapieerfolgskieferorthopädischer Behandlungen [Med. Diss.]: Heinrich-Heine-Universität Düsseldorf; 2001.
39. Miethe RR, Wronski C. What can be achieved with removable orthodontic appliances? Fortschr Kieferorthop/J Orofac Orthop 2009;70(3):185-99.
40. Riedmann T, Berg R. Retrospective evaluation of the outcome of orthodontic treatment in adults. Fortschr Kieferorthop/J Orofac Orthop 1999;60(2):108-23.
41. von Bremen J, Pancherz H. Efficiency of early and late Class II Division 1 treatment. Am J Orthod Dentofacial Orthop 2002;121(1):31-7.
42. von Bremen J, Bock N, Ruf S. Is Herbst-multibracket appliance treatment more efficient in adolescents than in adults? Angle Orthod 2009;79(1):173-7.
43. Prytz Berset G, Lund Eilertsen IM, Lagerstrom L, Espeland L, Stenvik A. Outcome of a scheme for specialist orthodontic care. Swed Dent J 2000;24(1-2):39-48.
44. Jeremiah HG, Cousley RR, Newton T, Abela S. Treatment time and occlusal outcome of orthognathic therapy in the East of England region. J Orthod 2012;39(3):206-11.
45. Pinzan-Vercelino CR, Janson G, Pinzan A, et al. Comparative efficiency of Class II malocclusion treatment with the pendulum appliance or two maxillary premolar extractions and edgewise appliances. Eur J Orthod 2009;31(3):333-40.
46. Richmond S, Andrews M. Orthodontic treatment standards in Norway. Eur J Orthod 1993;15(1):7-15.
47. Radznic D. Effectiveness of community-based salaried orthodontic services provided in England and Wales. J Orthod 2002;29(2):119-23.
48. Hönn M, Schneider C, Dietz K, Godt A, Göz G. Die Behandlung von Klasse-II-Patienten mit abnehmbaren Platten und funktionskieferorthopädischen Geräten. Fortschr Kieferorthop/J Orofac Orthop 2006;67(4):272-88.
49. Richmond S, Shaw WC, Stephens CD, et al. Orthodontics in the general dental service of England and Wales: a critical assessment of standards. Br Dent J 1993;174(9):315-29.
50. Richmond S, Andrews M, Roberts CT. The provision of orthodontic care in the general dental services of England and Wales: extraction patterns, treatment duration, appliance types and standards. Br J Orthod 1993;20(4):345-50.
51. Turbill EA, Richmond S, Wright JL. A critical assessment of orthodontic standards in England and Wales (1990-1991) in relation to changes in prior approval. Br J Orthod 1996;23(3):221-8.
52. Fox NA. The first 100 cases: a personal audit of orthodontic treatment assessed by the PAR (peer assessment rating) index. Br Dent J 1993;174(8):290-7.
53. Fox NA, Richmond S, Wright JL, Daniels CP. Factors affecting the outcome of orthodontic treatment within the general dental service. Br J Orthod 1997;24(3):217-21.
54. O'Brien KD, Shaw WC, Roberts CT. The use of occlusal indices in assessing the provision of orthodontic treatment by the hospital orthodontic service of England and Wales. Br J Orthod 1993;20(1):25-35.
55. Barton S, Cook PA. Predicting functional appliance treatment outcome in Class II malocclusions--a review. Am J Orthod Dentofacial Orthop 1997;112(3):282-6.
56. Murray AM. Discontinuation of orthodontic treatment: a study of the contributing factors. Br J Orthod 1989;16(1):1-7.
57. O'Brien K, Wright J, Conboy F, et al. Effectiveness of treatment for Class II malocclusion with the Herbst or twin-block appliances: a randomized, controlled trial. Am J Orthod Dentofacial Orthop 2003;124(2):128-37.
58. Ghafari J, Shofer FS, Jacobsson-Hunt U, Markowitz DL, Laster LL. Headgear versus function regulator in the early treatment of Class II, division 1 malocclusion: a randomized clinical trial. Am J Orthod Dentofacial Orthop 1998;113(1):51-61.
59. Sahm G, Bartsch A, Witt E. Micro-electronic monitoring of functional appliance wear. Eur J Orthod 1990;12(3):297-301.
60. Schott TC, Ludwig B. Microelectronic wear-time documentation of removable orthodontic devices detects heterogeneous wear behavior and individualizes treatment planning. Am J Orthod Dentofacial Orthop 2014;146(2):155-60.
61. Petren S, Bjerklund K, Marke LA, Bondemark L. Early correction of posterior crossbite--a cost-minimization analysis. Eur J Orthod 2013;35(1):14-21.
- Meyer-Gutknecht H, Fritz U, Schott TC. Methods to evaluate compliance of patients with removable appliances-survey results. Fortschr Kieferorthop/J Orofac Orthop 2014;75(2):144-53.

62. Schrömer R. Kieferorthopädie unter den Gesichtspunkten eingeschränkter Ressourcen. Zahnärztlicher Gesundheitsdienst 2000(01):5-6.
63. O'Brien K, Wright J, Conboy F, et al. Effectiveness of early orthodontic treatment with the Twin-block appliance: a multicenter, randomized, controlled trial. Part 1: Dental and skeletal effects. Am J Orthod Dentofacial Orthop 2003;124(3):234-43; quiz 339.
64. Demisch A. Langzeitbeobachtungen über die Stabilität der Okklusion nach Distalbisstherapie mit dem Berner Aktivator. Schweiz Monatsschr Zahnheilkd 1980;90(9):867-80.
65. Teuscher U. A growth-related concept for skeletal class II treatment. Am J Orthod 1978;74(3):258-75.
66. Jarvinen S, Widstrom E. Determinants of costs of orthodontic treatment in the Finnish public health service. Swed Dent J 2002;26(1):41-9.
67. Jarvinen S, Widstrom E, Raitio M. Factors affecting the duration of orthodontic treatment in children. A retrospective study. Swed Dent J 2004;28(2):93-100.
68. King GJ, McGorray SP, Wheeler TT, Dolce C, Taylor M. Comparison of peer assessment ratings (PAR) from 1-phase and 2-phase treatment protocols for Class II malocclusions. Am J Orthod Dentofacial Orthop 2003;123(5):489-96.
69. Tulloch JF, Proffit WR, Phillips C. Outcomes in a 2-phase randomized clinical trial of early Class II treatment. Am J Orthod Dentofacial Orthop 2004;125(6):657-67.
70. Gianelly AA. One-phase versus two-phase treatment. Am J Orthod Dentofacial Orthop 1995;108(5):556-9.
71. Johnston LE. Growing jaws for fun and profit: a modest proposal. In: McNamara JA, editor. Growth Modification: what works, what doesn't, and why. Ann Arbor: Center for Human Growth and Development, University of Michigan; 1999. p. 63-86.
72. Turbill EA, Richmond S, Wright JL. The time-factor in orthodontics: what influences the duration of treatments in National Health Service practices? Community Dent Oral Epidemiol 2001;29(1):62-72.